

DENSITY INVARIANCE OF CERTAIN OPERATIONAL QUANTITIES OF BOUNDED LINEAR OPERATORS IN NORMED SPACES

MICHAEL K. KINYON

ABSTRACT. Based on ideas of R.W. Cross, a simplified proof is presented of the density invariance of certain operational quantities associated with bounded linear operators in normed vector spaces.

Let X, Y denote normed linear spaces and let $T : X \rightarrow Y$ be a bounded linear operator. Let $\mathcal{I}(X)$ denote the collection of infinite dimensional subspaces of X . For any subspace M of X , let $S_M = \{m \in M : \|m\| = 1\}$ and let $T|_M$ denote the restriction of T to M . We introduce the following quantities:

$$\begin{aligned}\Gamma(T) &= \inf_{M \in \mathcal{I}(X)} \|T|_M\| \\ \Delta(T) &= \sup_{M \in \mathcal{I}(X)} \Gamma(T|_M) \\ \tau(T) &= \sup_{M \in \mathcal{I}(X)} \inf_{m \in S_M} \|Tm\| \\ \nabla(T) &= \inf_{M \in \mathcal{I}(X)} \tau(T|_M)\end{aligned}$$

These quantities were originally introduced in [6], [7], and [8] for bounded linear operators. They were extended to unbounded linear operators in [1] and [2], and to linear relations in [4]. See these papers for applications of these quantities to the study of bounded and unbounded linear operators and relations.

A linear subspace E of X is said to be a *core* of T if the graph of $T|_E$ is dense in the graph of T . A quantity f is said to be *densely invariant* if $f(T|_E) = f(T)$ whenever E is a core of T .

Our main result is the density invariance of the quantities Γ , Δ , τ , and ∇ . For Γ , Δ , and τ , this result first appeared in [2]; for ∇ , the first appearance is in [3]. The purpose of this note is to give a simplified proof. While the underlying ideas are the same as those in [2] and [3], the present proof has the following advantages: (1) the role played by the estimates is clearer, which considerably shortens the proof, and (2) the proof of the density invariance of Δ is direct and more obviously parallel to the proof of the density invariance of ∇ .

Our proof is for the case of bounded operators, but the case of unbounded operators reduces to this case [2], and the case of linear relations reduces to the case of unbounded operators [3].

We require the following [5, Lemma IV.2.8(ii)]:

The original version of this paper was written in the summer of 1995 while Ronald W. Cross was visiting the author at Indiana University South Bend. The author wishes to thank Prof. Cross for sharing his insights into linear operators and relations.

Lemma 1. *Let E be a dense linear subspace of X and let M be a closed, finite codimensional subspace of X . Then $E \cap M$ is dense in M .*

Theorem 1. *The quantities Γ , Δ , τ , and ∇ are densely invariant.*

Proof. Let E be a core of T . The case $\dim X < \infty$ is trivial, so we assume that X is infinite dimensional. For $M \in \mathcal{I}(X)$, we construct sequences $\{m_n\} \subset M$ and $\{x'_n\} \subset X'$ (here X' is the dual of X) as follows. Choose $m_1 \in S_M$ arbitrarily and choose x'_1 so that $\|x'_1\| = x'_1 m_1 = 1$. Suppose that $\{m_1, \dots, m_{n-1}\}$ and $\{x'_1, \dots, x'_{n-1}\}$ have been chosen. Choose m_n and x'_n so that

$$(1) \quad x'_i m_n = 0, \quad 1 \leq i < n,$$

$$(2) \quad \|m_n\| = \|x'_n\| = x'_n m_n = 1.$$

Let $W = \text{span}\{m_1, m_2, \dots\}$. It can be verified using (1) and (2) that the sequence $\{m_n\}$ is linearly independent. Thus $W \in \mathcal{I}(X)$. For an arbitrary element $w = \sum_1^n a_i m_i$, we shall verify that

$$(3) \quad |a_i| \leq 2^{i-1} \|w\|, \quad 1 \leq i \leq n.$$

Now $a_1 = x'_1 w$ by (1), so $|a_1| \leq \|x'_1\| \|w\| = \|w\|$. Suppose that (3) has been verified up to $i = j$. From (1) and (2), $x'_{j+1} w = \sum_1^j a_i x'_{j+1} m_i + a_{j+1}$. By the induction hypothesis,

$$|a_{j+1}| \leq |x'_{j+1} w| + \sum_1^j |a_i| |x'_{j+1} m_i| \leq \|w\| + \sum_1^j 2^{i-1} \|w\| \leq 2^j \|w\|$$

as required.

Let ϵ be given with $0 < \epsilon < 1$. For each n , the intersection of the subspace E with the intersection N of the kernels of the bounded linear functionals $x'_1, \dots, x'_n$ is dense in N by the lemma, and hence there exists a sequence $\{z_n\}$ in E such that

$$(4) \quad \|z_n - m_n\| \leq 2^{1-2n} \epsilon \min\{1, c/\|T\|\}$$

and

$$(5) \quad x'_i z_n = 0, \quad 1 \leq i < n$$

where $c > 0$ is an arbitrary constant.

Let $L = \text{span}\{z_1, z_2, \dots\}$. The linear independence of the sequence $\{z_n\}$ follows readily from (4), (5), and the linear independence of $\{m_n\}$. Thus $L \in \mathcal{I}(E)$.

Define a linear bijection $A : L \rightarrow W$ by $Az_i = m_i$. For $z = \sum_1^n a_i z_i \in L$, we have

$$(6) \quad \begin{aligned} \|z - Az\| &\leq \sum_1^n |a_i| \|z_i - Az_i\| \leq \sum_1^n 2^{i-1} \|Az\| \|z_i - Az_i\| \\ &\leq \epsilon \sum_1^n 2^{i-1} 2^{1-2i} \min\{1, c/\|T\|\} \|Az\| < \epsilon \min\{1, c/\|T\|\} \|Az\| \end{aligned}$$

by (3) and (4). Consequently

$$\begin{aligned}
 (1 - \epsilon)\|Az\| &\leq (1 - \epsilon \min\{1, c/\|T\|\})\|Az\| \\
 &< \|Az\| - \|z - Az\| \\
 &\leq \|z\| \\
 (7) \quad &\leq \|Az\| + \|z - Az\| \\
 &< (1 + \epsilon \min\{1, c/\|T\|\})\|Az\| \\
 &\leq (1 + \epsilon)\|Az\|.
 \end{aligned}$$

For $z \in L$, we have

$$\begin{aligned}
 \|Tz\| &\geq \|TAz\| - \|T(z - Az)\| \\
 &\geq \|TAz\| - \|T\|\|z - Az\| \\
 &> \|TAz\| - \|T\|\epsilon \min\{1, c/\|T\|\}\|Az\| \\
 &\geq \|TAz\| - \epsilon c\|Az\|.
 \end{aligned}$$

by (6). Thus using (7),

$$(8) \quad \frac{\|Tz\|}{\|z\|} > \left(\frac{\|TAz\|}{\|Az\|} - \epsilon c \right) (1 + \epsilon)^{-1}.$$

Similarly, we have for $z \in L$,

$$\begin{aligned}
 \|Tz\| &\leq \|TAz\| + \|T(z - Az)\| \\
 &\leq \|TAz\| + \|T\|\|z - Az\| \\
 &< \|TAz\| + \|T\|\epsilon \min\{1, c/\|T\|\}\|Az\| \\
 &\leq \|TAz\| + \epsilon c\|Az\|
 \end{aligned}$$

by (6). Thus using (7),

$$(9) \quad \frac{\|Tz\|}{\|z\|} < \left(\frac{\|TAz\|}{\|Az\|} + \epsilon c \right) (1 - \epsilon)^{-1}.$$

(a) The quantity Γ : We clearly have $\Gamma(T) \leq \Gamma(T|_E)$. Suppose that $\Gamma(T) < \Gamma(T|_E) - \delta$ where $\delta > 0$. Set $c = \Gamma(T|_E) - \delta$. Choose $M \in \mathcal{I}(X)$ such that

$$(10) \quad \|T|_M\| < c.$$

Then from (9) and (10)

$$\frac{\|Tz\|}{\|z\|} < \frac{1 + \epsilon}{1 - \epsilon} c$$

for $z \in L$. Hence

$$\Gamma(T|_E) = \inf_{V \in \mathcal{I}(E)} \|T|_V\| \leq \|T|_L\| \leq \frac{1 + \epsilon}{1 - \epsilon} c.$$

Since ϵ is arbitrary, $\Gamma(T|_E) \leq c$, a contradiction.

(b) The quantity τ : We clearly have $\tau(T) \geq \tau(T|_E)$. Suppose that $\tau(T) > \tau(T|_E) + \delta$ where $\delta > 0$. Set $c = \tau(T|_E) + \delta$. Choose $M \in \mathcal{I}(X)$ such that

$$(11) \quad \frac{\|Tm\|}{\|m\|} > c \text{ for } m \in M.$$

Then from (8) and (11)

$$\frac{\|Tz\|}{\|z\|} > \frac{1-\epsilon}{1+\epsilon}c$$

for $z \in L$. Hence

$$\tau(T|_E) \geq \inf_{z \in L} \frac{\|Tz\|}{\|z\|} \geq \frac{1-\epsilon}{1+\epsilon}c.$$

Since ϵ is arbitrary, $\tau(T|_E) \geq c$, a contradiction.

(c) The quantity Δ : We clearly have $\Delta(T|_E) \leq \Delta(T)$. Let $\delta > 0$ be given and choose $M_0 \in \mathcal{I}(X)$ such that $\Gamma(T|_{M_0}) > \Delta(T) - \delta$. Set $c = \Delta(T) - \delta$. Then if $M \in \mathcal{I}(M_0)$,

$$(12) \quad \|T|_M\| > c.$$

From (8) we have

$$\|T|_V\| \geq (\|T|_{AV}\| - \epsilon c)(1 + \epsilon)^{-1}$$

for $V \in \mathcal{I}(L)$. Since $AV \in \mathcal{I}(M) \subset \mathcal{I}(M_0)$, it follows from (12) that

$$\|T|_V\| \geq \frac{1-\epsilon}{1+\epsilon}c$$

for $V \in \mathcal{I}(L)$. Thus

$$\Gamma(T|_L) = \inf_{V \in \mathcal{I}(L)} \|T|_V\| \geq \frac{1-\epsilon}{1+\epsilon}c.$$

Taking the supremum of the left side over $L \in \mathcal{I}(E)$, we have

$$\Delta(T|_E) \geq \frac{1-\epsilon}{1+\epsilon}c = \frac{1-\epsilon}{1+\epsilon}(\Delta(T) - \delta).$$

Since ϵ and δ are arbitrary, it follows that $\Delta(T|_E) \geq \Delta(T)$.

(d) The quantity ∇ : We clearly have $\nabla(T|_E) \geq \nabla(T)$. Let $\delta > 0$ be given and choose $M_0 \in \mathcal{I}(X)$ such that $\tau(T|_{M_0}) < \nabla(T) + \delta$. Set $c = \nabla(T) + \delta$. Then if $M \in \mathcal{I}(M_0)$,

$$(13) \quad \inf_{m \in S_M} \|Tm\| < c.$$

From (9) we have

$$\inf_{z \in V} \frac{\|Tz\|}{\|z\|} \leq \left(\inf_{Az \in AV} \frac{\|TAz\|}{\|Az\|} + \epsilon c \right) (1 - \epsilon)^{-1}$$

for $V \in \mathcal{I}(L)$. Since $AV \in \mathcal{I}(M) \subset \mathcal{I}(M_0)$, it follows from (13) that

$$\inf_{z \in V} \frac{\|Tz\|}{\|z\|} \leq \frac{1+\epsilon}{1-\epsilon}c$$

for $V \in \mathcal{I}(L)$. Thus

$$\tau(T|_L) = \sup_{V \in \mathcal{I}(L)} \inf_{z \in V} \frac{\|Tz\|}{\|z\|} \leq \frac{1+\epsilon}{1-\epsilon}c.$$

Taking the infimum of the left side over $L \in \mathcal{I}(E)$, we have

$$\nabla(T|_E) \leq \frac{1+\epsilon}{1-\epsilon}c = \frac{1+\epsilon}{1-\epsilon}(\nabla(T) + \delta).$$

Since ϵ and δ are arbitrary, it follows that $\nabla(T|_E) \leq \nabla(T)$. □

REFERENCES

- [1] R.W. Cross, Properties of some norm related functions of unbounded linear operators, *Math. Z.* **199** (1988), 285-302.
- [2] R.W. Cross, On certain densely invariant quantities of linear operators, *Math. Nachr.* 178 (1996), 103-116.
- [3] R.W. Cross, *Multivalued Linear Operators*, Marcel Dekker, Pure and Appl. Math. **213**, 1998.
- [4] R.W. Cross and P. Pillay, Ubiquitous properties of certain operational quantities of linear relations in normed spaces, *Quaestiones Mathematicae*, **17** (1994), 487-498.
- [5] S. Goldberg, *Unbounded Linear Operators: Theory and Applications*, Dover Publications Inc., 1966.
- [6] A. Lebow and M. Schechter, Semigroups of operators and measures of noncompactness, *J. Funct. Anal.* **7** (1971), 1-26.
- [7] M. Schechter, Quantities related to strictly singular operators, *Indiana Univ. Math. J.* **21** (1972), 1061-1071.
- [8] A. Sedaev, The structure of certain linear operators, *Mat. Issled* **5** (1970), 166-175. (Russian)

DEPARTMENT OF MATHEMATICS, WESTERN MICHIGAN UNIVERSITY, KALAMAZOO, MI USA
E-mail address: mkinyon@wmich.edu